CANADIAN JOURNAL OF FOREST RESEARCH

JOURNAL CANADIEN DE LA RECHERCHE FORESTIÈRE

VOLUME 8, 1978

Author Index/Index des auteurs

Under each name are listed, in chronological order of publication, all articles in which an author has participated/Chaque notice indique, dans l'ordre chronologique de publication, tous les articles auxquels l'auteur a participé

	Baradat, Ph.	
nes on	Variabilité et hérédité de l'angle du fil du bois mes	uré à
	l'aide d'un traceur radioactif chez le pin maritime et l	e pin
	laricio de Calabre	
306	M. ARBEZ, Ph. BARADAT, Y. BIROT, P. AZOEUF et	
	R. HOSLIN	280
	Rarney P I	
spruce		13374
		LANC
290		36
	NODERI OCHEENINER	50
	Racham IT	
tion de		as mon
non de		spen
111		181
111	K. HUNI, J. I. DASHAM, allu J.A. KEMPERMAN	101
	Poster S.M.	
esuré à		121
le pin	SHELAGH WI. DAXIER AND MELVIN G.R. CANNELL	121
	Bédard, J.	
280		of ar
	early seral wintering site in Gaspé Peninsula, Quebec	
	J. BÉDARD, M. CRÊTE, and E. AUDY	407
Clado-	Beineke, W.F.	
168		346
	Rirot V	
of an		uré i
	laricio de Calabre	1
	M. ARBEZ, PH. BARADAT, Y. BIROT, P. AZOEUF CL	
	R. Hoslin	280
esuré à	Planchard P ()	
t le pin		nla
		pic
		90
280	NOBERT O. BLANCHARD	×
	Blanchard B O	
	306 spruce 290 tion de 111 esuré à t le pin 280 Clado- 168 of an 407	Notes on Variabilité et hérédité de l'angle du fil du bois mes l'aide d'un traceur radioactif chez le pin maritime et l'aricio de Calabre 306 M. Arbez, Ph. Baradat, Y. Birot, P. Azoeuf et R. Hoslin Spruce Barney, R.J. Biomass distribution and crown characteristics in Alaskan Picea mariana ecosystems RICHARD J. Barney, Keith Van Cleve, and Robert Schlentner Basham, J.T. Kraft-pulping evaluation of decayed trembling a (Populus tremuloides) from Ontario 111 K. Hunt, J.T. Basham, and J.A. Kemperman Baxter, S.M. Branch development on leaders of Picea sitchensis Shelagh M. Baxter and Melvin G.R. Cannell. Bédard, J. Short-term influence of moose upon woody plants early seral wintering site in Gaspé Peninsula, Quebec J. Bédard, M. Crête, and E. Audy Clado-Beineke, W.F. Irrigation protects black walnut from frost damage Walter F. Beineke Walter F. Beineke Birot, Y. Variabilité et hérédité de l'angle du fil du bois mes l'aide d'un traceur radioactif chez le pin maritime et laricio de Calabre M. Arbez, Ph. Baradat, Y. Birot, P. Azoeuf et R. Hoslin Blanchard, R.O. Electrical resistance related to phloem width in red ma Jonathan K. Carter and Robert O. Blanchard

Within-stand seedling dispersal for isolated Pinus strobus

within hardwood stands

V.A. RUDIS, A.R. EK, and J.W. BALSIGER

Effects of soil-applied potassium on cation distribution

ROBERT O. BLANCHARD, DESMOND SMITH, ALEX L. SHIGO,

228

around wounds in red maple

and L.O. SAFFORD

Blanchette, R.A.	Cahill, M.J.
A modified technique for the determination of fungal mass	A pattern of forest types on ribbed moraines in eastern
in decayed wood	Newfoundland
S. GURUSIDDAIAH, ROBERT A. BLANCHETTE, and C. GARDNER SHAW 486	B.B. Delaney and M.J. Cahill 116
C. GARDINER SHAW	Cannell, M.G.R.
Bloomberg, W.J.	Branch development on leaders of Picea sitchensis
Heatsum-emergence relationship in Douglas-fir seedlings	SHELAGH M. BAXTER and MELVIN G.R. CANNELL 121
W.J. Bloomberg 23	
	Cannell, M.G.R.
Bonnet-Masimbert, M.	Spatial arrangement of lateral buds at the time that they form on leaders of <i>Picea</i> and <i>Larix</i>
Induction florale sur pousses d'août chez le Douglas (Pseudotsuga menziesii)	M.G.R. CANNELL and K.C. Bowler 129
MARC BONNET-MASIMBERT et René Lanares 247	
	Cannell, M.G.R.
Botkin, D.B.	Phyllotactic arrangements of needles on elongating conifer
Predicting the effects of different harvesting regimes on	shoots: a computer simulation M.G.R. CANNELL and K.C. BOWLER 138
forest floor dynamics in northern hardwoods	M.O.R. CANNELL and R.C. BOWLER
JOHN D. ABER, DANIEL B. BOTKIN, and JERRY M. MELILLO 306	Carter, J.K.
JERRY IVI. IVIELLELO	Electrical resistance related to phloem width in red maple
Bowdery, L.	JONATHAN K. CARTER and
Effects of nursery seedbed density and topdressing fertil-	ROBERT O. BLANCHARD 90
ization on survival and growth of 3+0 red pine	
R.E. MULLIN and L. BOWDERY 30	Christersson, L. Short-term temperature variation in needles of <i>Pinus</i>
	silvestris L.
Bowler, K.C.	Lars Christersson and Roland Sandstedt 480
Spatial arrangement of lateral buds at the time that they	
form on leaders of <i>Picea</i> and <i>Larix</i> M.G.R. CANNELL and K.C. BOWLER 129	Cown, D.J.
M.O.R. CANNELL and R.C. BOWLER	Comparison of annual ring density profiles in hardwoods
Bowler, K.C.	and softwoods by X-ray densitometry
Phyllotactic arrangements of needles on elongating conifer	D.J. Cown and M.L. Parker 442
shoots: a computer simulation	Crête, M.
M.G.R. CANNELL and K.C. BOWLER 138	Short-term influence of moose upon woody plants of an
	early seral wintering site in Gaspé Peninsula, Quebec
Bradbury, I.K.	J. BÉDARD, M. CRÊTE, and E. AUDY 407
Dry matter accumulation by <i>Picea sitchensis</i> seedlings during winter	
IAN K. BRADBURY and D.C. MALCOLM 207	D'Aoust, A.L.
	Influence de l'irradiation sur la croissance de plants d'épinette noire en contenants placés dans deux enceintes
Britnell, W.E.	de culture
Comparison of malt agar with malt agar plus ortho-	André L. D'Aoust 316
phenylphenol for isolating Armillaria mellea and other fungi	
from conifer roots R.D. Whitney, D.T. Myren, and W.E. Britnell. 348	Delaney, B.B.
N.D. WHITSET, D. I. WITKEN, allU W.E. DKIINELL 346	A pattern of forest types on ribbed moraines in eastern
Burdett, A.N.	Newfoundland B.B. Delaney and M.J. Cahill 116
Control of root morphogenesis for improved mechanical	D.D. DELANET and M.J. CAHLL
stability in container-grown lodgepole pine	Denne, M.P.
A.N. BURDETT 483	Variation of fibre length within trees of Fraxinus excelsior
	M.P. Denne and V. Whitbread 253
Burkhart, H.E.	
Allocating inventory resources for multiple-use planning	Densmore, R.
HAROLD E. BURKHART, R. DEAN STUCK, WILLIAM A. LEUSCHNER, and	Rooting potential of Alaskan willow cuttings
MARION R. REYNOLDS 100	ROSEANN DENSMORE and JOHN C. ZASADA 477
100	Dickmann, D.I.
Byron, R.N.	Marked differences among poplar clones in winter browsing
Community stability and forest policy in British Columbia	damage by cottontail rabbits
R.N. Byron 61	DONALD I. DICKMANN 351


Dixon, M.A. Electrical resistance measurements of water potential in	Gurusiddaiah, S. A modified technique for the determination of fungal mass
avocado and white spruce M.A. Dixon, R.G. Thompson, and D.S. Fensom 73	in decayed wood S. Gurusiddaiah, Robert A. Blanchette, and
D 1.1 W.S	C. GARDNER SHAW 486
Drechsler, H.D. The impact of composite sampling and other data aggre-	Hardy, Y.J.
gation procedures on pollution detection in the pulp and paper industry	An assay of sequential application of Fenitrothion and Bacillus thuringiensis for an integrated control of the spruce
HERBERT D. DRECHSLER and PETER N. NEMETZ 328	budworm
Durzan, D.J.	W.A. SMIRNOFF and Y.J. HARDY 300
Effect of fertilization of balsam fir trees on spruce budworm	Hare, R.C.
nutrition and development	Effect of shoot girdling and season on rooting of slash pine
G.G. Shaw, C.H.A. LITTLE, and D.J. DURZAN 364	cuttings Robert C. Hare
Ek, A.R.	
Within-stand seedling dispersal for isolated <i>Pinus strobus</i> within hardwood stands	Harvey, A.E.
V.A. Rudis, A.R. Ek, and J.W. Balsiger 10	N ₂ fixation associated with wood decayed by some common
	fugi in western Montana Michael J. Larsen, Martin F. Jurgensen, and
Ek, A.R.	ALAN E. HARVEY 341
Reduced growth, cull, and mortality of jack pine associated	
with sweetfern rust cankers HENRY L. GROSS, ROBERT F. PATTON, and	Hawkes, B.C.
ALAN R. EK 47	Distribution and chemistry of fine roots in a white spruce - subalpine fir stand in British Columbia: implications for
Fasehun, F.E.	Management J.P. Kimmins and B.C. Hawkes 265
Effect of irradiance on growth and photosynthesis of	JA . REMAINS WIN D.C. ITAWALD
Populus × euramericana clones F.E. Fasehun 94	Hodgson, M.J.
F.E. FASEHUN 94	Location-allocation models for one-strike initial attack of
Fensom, D.S.	forest fires by airtankers
Electrical resistance measurements of water potential in	M.J. Hodgson and R.G. Newstead 145
avocado and white spruce	Hoslin, R.
M.A. Dixon, R.G. Thompson, and D.S. Fensom 73	Variabilité et hérédité de l'angle du fil du bois mesuré à
Fisher, R.F.	l'aide d'un traceur radioactif chez le pin maritime et le pin
Allelopathic effects of goldenrod and aster on young sugar	laricio de Calabre
maple	M. Arbez, Ph. Baradat, Y. Birot, P. Azoeuf et R. Hoslin 280
R.F. FISHER, R.A. WOODS, and M.R. GLAVICIC 1	TO TAXABLE
Clarieia M.B.	Hunt, K.
Glavicic, M.R. Allelopathic effects of goldenrod and aster on young sugar	Kraft-pulping evaluation of decayed trembling aspen
maple	(Populus tremuloides) from Ontario
R.F. FISHER, R.A. WOODS, and M.R. GLAVICIC 1	K. Hunt, J.T. Basham, and J.A. Kemperman 181
Golding, D.L.	Johnson, A.L.S.
Snow accumulation and melt in small forest openings in	Stump colonization and spread of Fomes annosus 5 years
Alberta	after thinning
D.L. Golding and R.H. Swanson 380	D.J. Morrison and A.L.S. Johnson 177
Grier, C.C.	Jurgensen, M.F.
A Tsuga heterophylla - Picea sitchensis ecosystem of coastal	
Oregon: decomposition and nutrient balances of fallen logs	fugi in western Montana Michael J. Larsen, Martin F. Jurgensen, and
Charles C. Grier 198	ALAN E. HARVEY 341
Gross, H.L.	
Reduced growth, cull, and mortality of jack pine associated	
with sweetfern rust cankers	Southern pine stem form defined through principal compo-
HENRY L. GROSS, ROBERT F. PATTON, and ALAN R. EK 47	nent analysis C.J. Liu and T.D. Keister 188
ALAN IV. FV	C.J. LIU dild 1.D. REISTER 100

Kemperman, J.A. Kraft-pulping evaluation of decayed trembling (Populus tremuloides) from Ontario K. Hunt, J.T. Basham, and J.A. Kemperman	aspen	Lotan, J.E. Variation in lodgepole pine (<i>Pinus contorta</i> var. <i>latifolia</i>): greenhouse response of wind pollinated families from five populations to day length and temperature—soil
		DAVID A. PERRY and JAMES E. LOTAN 81
Kimmins, J.P.		
Distribution and chemistry of fine roots in a white spi subalpine fir stand in British Columbia: implication management		Mahendrappa, M.K. Changes in the organic layers under a black spruce stand
J.P. KIMMINS and B.C. HAWKES	265	fertilized with urea and triple superphosphate M.K. Mahendrappa 237
Kozak, A. The potential of Weibull-type functions as flexible greaters.	rowth	Malcolm, D.C. Dry matter accumulation by <i>Picea sitchensis</i> seedlings
R.C. YANG, A. KOZAK, and J.H.G. SMITH	424	during winter IAN K. BRADBURY and D.C. MALCOLM 207
		TAN K. DRADBURT and D.C. WALLOLM
Lanares, R.		Malia, M.E.
Induction florale sur pousses d'août chez le De (Pseudotsuga menziesii)	ouglas	Electrical resistance, physical characteristics, and cation
MARC BONNET-MASIMBERT et RENÉ LANARES	247	concentrations in xylem of sugar maple infected with Verticillium dahliae
Larsen, M.J.		MARGARET E. MALIA and TERRY A. TATTAR 322
N ₂ fixation associated with wood decayed by some cor	nmon	
fugi in western Montana		Martin, G.L.
MICHAEL J. LARSEN, MARTIN F. JURGENSEN, and	241	Illustration of nursery stock distribution by computer graphics
ALAN E. HARVEY	341	D.T. LESTER and G.L. MARTIN 261
Larson, M.M.		
Effects of late-season defoliation and dark periods on	initial	McCaughey, J.H.
growth of planted northern red oak seedlings		Estimation of net radiation for a coniferous forest, and the
M.M. Larson	67	effects of logging on net radiation and the reflection coefficient
Leaphart, C.D.	0	J.H. McCaughey 450
A system using aerial photography to estimate area o	f root	
disease centers in forests RALPH E. WILLIAMS and C.D. LEAPHART	214	McCaughey, J.H. Energy balance and evapotranspiration estimates for a
NABITE WILLIAMS and C.D. BEATTAKT	214	mature coniferous forest
Lester, D.T.		J.H. McCaughey 456
Illustration of nursery stock distribution by com	puter	
graphics D.T. Lester and G.L. Martin	2/1	McClenahen, J.R.
	261	Community changes in a deciduous forest exposed to air pollution
Leuschner, W.A. Allocating inventory resources for multiple-use planning	20	JAMES R. MCCLENAHEN 432
HAROLD E. BURKHART, R. DEAN STUCK,	1g	Column and
WILLIAM A. LEUSCHNER, and		Melillo, J.M.
Marion R. Reynolds	100	Predicting the effects of different harvesting regimes on forest floor dynamics in northern hardwoods JOHN D. ABER, DANIEL B. BOTKIN, and
Lheureux, C.		JERRY M. MELILLO 306
Réactions 'organographiques' du pin laricio sous l'acti	ion de	
l'acide cacodylique Christian Lheureux et Pierre Andre	111	Methyen, I.R.
	111	Discussion: Two recent articles on fire ecology C.E. VAN WAGNER and I.R. METHVEN 491
Little, C.H.A. Effect of fertilization of balsam fir trees on spruce bud	33173 00000	7/1
nutrition and development	worm	Moore, K.E.
G.G. Shaw, C.H.A. LITTLE, and D.J. DURZAN	364	Barrier-zone formation in wounded stems of sweetgum KATHLEEN E. MOORE 389
Liu, C.J.		
Southern pine stem form defined through principal co	ompo-	Morgenstern, E.K.
nent analysis C.J. Liu and T.D. Keister	100	Range-wide genetic variation of black spruce
C.J. LIU dilu 1.D. KEISTEK	188	E.K. MORGENSTERN 463

Morrison, D.J.	Perry, D.A.
Stump colonization and spread of Fomes annosus 5 years	Variation in lodgepole pine (Pinus contorta var. latifolia):
after thinning	greenhouse response of wind pollinated families from five
D.J. Morrison and A.L.S. Johnson 177	populations to day length and temperature-soil DAVID A. PERRY and JAMES E. LOTAN 81
	DAVID A. PERRY AND JAMES E. LOTAN 81
Mullin, R.E.	Piene, H.
Effects of nursery seedbed density and topdressing fertil-	Weight loss of litter and cellulose bags in a thinned white
ization on survival and growth of 3 + 0 red pine	spruce forest in interior Alaska
R.E. MULLIN and L. BOWDERY 30	HARALD PIENE and KEITH VAN CLEVE 42
Myren, D.T.	Piene, H.
Root-rotting fungi associated with mortality of conifer	Effects of increased spacing on carbon mineralization rates
saplings in northern Ontario	and temperature in a stand of young balsam fir
ROY D. WHITNEY and DONALD T. MYREN 17	HARALD PIENE 398
	Pomber, L.
Myren, D.T.	Phytotoxicity of fenitrothion to white pine seeds and
Comparison of malt agar with malt agar plus ortho-	seedlings
phenylphenol for isolating Armillaria mellea and other fungi from conifer roots	PEARL WEINBERGER, L. POMBER, and R. PRASAD 155
R.D. WHITNEY, D.T. MYREN, and W.E. BRITNELL 348	Pomber, L.
	Some toxic effects of fenitrothion on seed germination and
Nemetz, P.N.	early seedling growth of jack pine, spruce, and birches
The impact of composite sampling and other data aggre-	PEARL WEINBERGER, L. POMBER, and R. PRASAD 243
gation procedures on pollution detection in the pulp and	
paper industry	Prasad, R.
HERBERT D. DRECHSLER and PETER N. NEMETZ 328	Phytotoxicity of fenitrothion to white pine seeds and
	seedlings
Newstead, R.G.	PEARL WEINBERGER, L. POMBER, and R. PRASAD 155
Location-allocation models for one-strike initial attack of	Prasad, R.
forest fires by airtankers	Some toxic effects of fenitrothion on seed germination and
M.J. HODGSON and R.G. NEWSTEAD 145	early seedling growth of jack pine, spruce, and birches
	PEARL WEINBERGER, L. POMBER, and R. PRASAD 243
Owston, P.W.	
Container and root treatments affect growth and root form	Rencz, A.N.
of planted ponderosa pine	Biomass distribution in a subarctic Picea mariana - Clado-
PEYTON W. OWSTON and K.W. SEIDEL 232	nia alpestris woodland
	ANDREW N. RENCZ and ALLAN N.D. AUCLAIR 168
Parker, M.L.	
Comparison of annual ring density profiles in hardwoods	Reynolds, M.R.
and softwoods by X-ray densitometry	Allocating inventory resources for multiple-use planning
D.J. Cown and M.L. Parker 442	HAROLD E. BURKHART, R. DEAN STUCK, WILLIAM A. LEUSCHNER, and
	MARION R. REYNOLDS 100
Patton, R.F.	MARION R. RETROLDS
Reduced growth, cull, and mortality of jack pine associated	Riding, R.T.
with sweetfern rust cankers	The epicuticular waxes of Pinus strobus subjected to air
HENRY L. GROSS, ROBERT F. PATTON, and	pollutants
ALAN R. EK	K.E. Percy and R.T. Riding 474
Perala, D.A.	Rohde, R.A.
Biomass and nutrient distribution in aspen, pine, and spruce	Incidence and condition of vesicular-arbuscular mycorrhi-
stands on the same soil type in Minnesota	zae infections in the roots of sugar maple in relation to maple decline
DAVID H. ALBAN, DONALD A. PERALA, and	ROBERTA A. SPITKO, TERRY A. TATTAR, and
BRYCE E. SCHLAEGEL 290	RICHARD A. ROHDE 375
Percy, K.E.	Rudis, V.A.
The epicuticular waxes of Pinus strobus subjected to air	Within-stand seedling dispersal for isolated Pinus strobus
pollutants	within hardwood stands
K.E. Percy and R.T. RIDING 474	V.A. Rudis, A.R. Ek, and J.W. Balsiger 10

Safford, L.O.	Smith, J.H.G.
Effects of soil-applied potassium on cation distribution	The potential of Weibull-type functions as flexible growth
around wounds in red maple	D.C. V. vic. A. Konsak and I.H.C. States
ROBERT O. BLANCHARD, DESMOND SMITH, ALEX L. SHIGO, and L.O. SAFFORD 228	R.C. YANG, A. KOZAK, and J.H.G. SMITH 424
	Spitko, R.A.
Sandstedt, R.	Incidence and condition of vesicular-arbuscular mycorrhi
Short-term temperature variation in needles of <i>Pinus</i> silvestris L.	zae infections in the roots of sugar maple in relation to
LARS CHRISTERSSON and ROLAND SANDSTEDT 480	maple decline ROBERTA A. SPITKO, TERRY A. TATTAR, and
	RICHARD A. ROHDE 37:
Schlaegel, B.E.	
Biomass and nutrient distribution in aspen, pine, and spruce	Spomer, G.G.
stands on the same soil type in Minnesota	Comparisons of water potentials in bole and needle tissue
DAVID H. ALBAN, DONALD A. PERALA, and BRYCE E. SCHLAEGEL 290	of lodgepole pine
DRYCE E. SCHLAEGEL 290	GEORGE G. SPOMER and KAREL J. STOSZEK 43
Schlentner, R.	Stauffer, H.B.
Biomass distribution and crown characteristics in two	Aggregating points to fit Pielou's index of nonrandomness
Alaskan Picea mariana ecosystems RICHARD J. BARNEY, KEITH VAN CLEVE, and	HOWARD B. STAUFFER 35:
ROBERT SCHLENTNER 36	
TO THE STATE OF TH	Stoszek, K.J.
Seidel, K.W.	Comparisons of water potentials in bole and needle tissue
Container and root treatments affect growth and root form	of lodgepole pine
of planted ponderosa pine	GEORGE G. SPOMER and KAREL J. STOSZEK 43
PEYTON W. OWSTON and K.W. SEIDEL 232	
61 66	Stuck, R.D.
Shaw, C.G. A modified technique for the determination of fungal mass	Allocating inventory resources for multiple-use planning HAROLD E. BURKHART, R. DEAN STUCK,
in decayed wood	WILLIAM A. LEUSCHNER, and
S. GURUSIDDAIAH, ROBERT A. BLANCHETTE, and	MARION R. REYNOLDS 10
C. GARDNER SHAW 486	
	Swanson, R.H.
Shaw, G.G.	Snow accumulation and melt in small forest openings i
Effect of fertilization of balsam fir trees on spruce budworm nutrition and development	Alberta
G.G. Shaw, C.H.A. LITTLE, and D.J. DURZAN 364	D.L. GOLDING and R.H. SWANSON 38
	Sylvia, D.M.
Shigo, A.L.	Electrical resistance properties of tree tissues in canker
Discolouration and decay associated with paraformalde-	incited by Endothia parasitica and Nectria galligena
hyde-treated tapholes in sugar maple RUSSELL S. WALTERS and ALEX L. SHIGO 54	DAVID M. SYLVIA and TERRY A. TATTAR 16
NOSSELLO, WALTERS and PALEA L. SHOO	
Shigo, A.L.	Tattar, T.A.
Effects of soil-applied potassium on cation distribution	Electrical resistance properties of tree tissues in canker incited by <i>Endothia parasitica</i> and <i>Nectria galligena</i>
around wounds in red maple	David M. Sylvia and Terry A. Tattar 16
ROBERT O. BLANCHARD, DESMOND SMITH, ALEX L. SHIGO, and L.O. SAFFORD 228	DATID III. DILVIA GIO ILIANI II. IAITAK
and L.O. Safford 228	Tattar, T.A.
Smirnoff, W.A.	Electrical resistance, physical characteristics, and catio
An assay of sequential application of Fenitrothion and	concentrations in xylem of sugar maple infected wit
Bacillus thuringiensis for an integrated control of the spruce	Verticillium dahliae
budworm	MARGARET E. MALIA and TERRY A. TATTAR 32
W.A. SMIRNOFF and Y.J. HARDY 300	
Smith, D.	Tattar, T.A.
Effects of soil-applied potassium on cation distribution	Incidence and condition of vesicular-arbuscular mycorrh zae infections in the roots of sugar maple in relation t
around wounds in red maple	maple decline
ROBERT O. BLANCHARD, DESMOND SMITH, ALEX L. SHIGO,	ROBERTA A. SPITKO, TERRY A. TATTAR, and
and L.O. SAFFORD 228	RICHARD A. ROHDE 37

Tauer, C.G. Sweet fern rust resistance in jack pine seedlings: geograp	phic	Weinberger, P. Some toxic effects of fenitrothion on seed germination a	and
variation Charles G. Tauer	416	early seedling growth of jack pine, spruce, and birches PEARL WEINBERGER, L. POMBER, and R. PRASAD	243
CHARLES G. TAUER	410	FEARL WEINBERGER, L. FOMBER, AIR R. FRASAD	43
Thompson., R.G. Electrical resistance measurements of water potentia avocado and white spruce	l in	Whitbread, V. Variation of fibre length within trees of Fraxinus excelsion M.P. Denne and V. Whitbread	r 253
M.A. Dixon, R.G. THOMPSON, and D.S. FENSOM	73	Whitney, R.D. Root-rotting fungi associated with mortality of con-	ifer
Van Cleve, K. Biomass distribution and crown characteristics in	two	saplings in northern Ontario ROY D. WHITNEY and DONALD T. MYREN	17
Alaskan <i>Picea mariana</i> ecosystems Richard J. Barney, Keith Van Cleve, and Robert Schlentner	36	Whitney, R.D. Comparison of malt agar with malt agar plus ort phenylphenol for isolating <i>Armillaria mellea</i> and other fu	
Van Cleve, K. Weight loss of litter and cellulose bags in a thinned w	hite		348
spruce forest in interior Alaska Harald Piene and Keith Van Cleve	42	Williams, R.E. A system using aerial photography to estimate area of a disease centers in forests	
Van Wagner, C.E. Age-class distribution and the forest fire cycle		RALPH E. WILLIAMS and C.D. LEAPHART Woods, R.A.	214
C.E. VAN WAGNER	220	Allelopathic effects of goldenrod and aster on young su maple	ıga
Van Wagner, C.E. Discussion: Two recent articles on fire ecology		R.F. FISHER, R.A. WOODS, and M.R. GLAVICIC	
C.E. Van Wagner and I.R. Methven	491	Yang, R.C. The potential of Weibull-type functions as flexible grocurves	wtl
Walters, R.S. Discolouration and decay associated with paraforma	alda	R.C. YANG, A. KOZAK, and J.H.G. SMITH	424
hyde-treated tapholes in sugar maple		Zasada, J.C. Rooting potential of Alaskan willow cuttings	
RUSSELL S. WALTERS and ALEX L. SHIGO	54		47
Weinberger, P. Phytotoxicity of fenitrothion to white pine seeds	and	Ziemer, R.R. An apparatus to measure the crosscut shearing strengt	h o
seedlings	155	PODEST R ZIEMER	14



CANADIAN JOURNAL OF FOREST RESEARCH

JOURNAL CANADIEN DE LA RECHERCHE FORESTIÈRE

VOLUME 8, 1978

Subject Index/Index des matières

This is a rotated term index. The terms are arranged linearly, primary terms first, followed by secondary terms after the semicolon. Each primary term is used once to head an entry. Secondary terms are informative but not sufficiently specific to head entries. In an entry, a few terms by themselves may be ambiguous, but a reading of all the terms should convey the content of the paper. The index terms are followed by the volume number, inclusive page numbers, and author's names, in that order.

Il s'agit d'un index de termes permutés, présentés de façon linéaire: les termes principaux sont suivis, après le point-virgule, des termes secondaires. Chaque terme principal figure une fois au début d'une notice. Les termes secondaires sont utiles, mais trop peu précis pour servir de termes principaux. Il peut arriver que quelques termes d'une notice, pris isolément, soient obscurs, mais l'ensemble des termes doit préciser le contenu de l'article. Les termes de l'index sont suivis du numéro du volume, de la pagination complète et du nom de l'auteur ou des auteurs, dans cet ordre.

- Abies balsamea, Alces alces, moose, browse, regeneration; Quebec, Acer spicatum, Corylus cornuta. 8, 407–415 (Bédard et
- fertilization, budworm, Choristoneura fumiferana; New Brunswick, budworm nutrition. 8, 364–374 (Shaw et al.)
- ———, forest types, *Picea mariana, Betula papyrifera*; Newfoundland, ribbed moraines. **8**, 116–120 (Delaney and Cahill)
- spacing, mineralization, organic layers, temperature; Nova Scotia. 8, 398-406 (Piene)
- Abies lasiocarpa, Picea glauca, roots, nutrient losses; British Columbia, management, logging, fine-root biomass, mineral elements. 8, 265–279 (Kimmins and Hawkes)
- Acer rubrum, cation distribution, wounds; Maine, U.S.A., soil-applied potassium. 8, 228-231 (Blanchard et al.)
- -----, electrical resistance, phloem, Shigometer; field, Maine. 8, 90-93 (Carter and Blanchard)
- Acer saccharum, allelopathy; field, greenhouse, laboratory, fertilization, plantations. 8, 1–9 (Fisher et al.)
- ———, electrical resistance, Verticillium dahliae, cation concentration, decay; Massachusetts, U.S.A. 8, 322–327 (Malia and Tattar)
- ———, maple syrup, wound reaction, decay; northeastern United States, succession of microorganisms, paraformaldehyde. 8, 54–60 (Walters and Shigo)
- mycorrhizae; Massachusetts, maple decline. 8, 375-379 (Spitko et al.)
- ——, pollution; Ohio River Valley, U.S.A., airborne, seven stands, richness and deversity. **8**, 432–438 (McClenahen)
- age-class, fire; fire cycle, fire history. 8, 220-227 (Van Wagner)
- aggregation, Pielou's index; nonrandom spatial patterns. 8, 355-363 (Stauffer)
- Alces alces, Abies balsamea, moose, browse, regeneration; Quebec, Acer spicatum, Corylus cornuta. 8, 407-415 (Bédard et al.)
- allelopathy, Acer saccharum; field, greenhouse, laboratory, fertilization, plantations. 8, 1-9 (Fisher et al.)
- annual ring, X-ray densitometry; radial and transverse sections. 8, 442-449 (Cown and Parker)
- apical dominance, bud, morphogenesis, *Picea sitchensis*, plastochrone, primordia; Great Britain, cell dimensions. 8, 121–128 (Baxter and Cannell)
- Armillaria mellea, infection centers, Fomes annosus, Pseudotsuga menziesii, Tsuga heterophylla; stumps, competition. 8, 177–180 (Morrison and Johnson)
 - , isolation; selective media. 8, 348-350 (Whitney et al.)
- , Scytinostroma galactinum, Coniophora puteana, Pinus banksiana, Pinus resinosa, Abies balsamea, root rotting; conifer saplings, northern Ontario. 8, 17–22 (Whitney and Myren)

Bacillus thuringiensis, spruce budworm, Choristoneura fumiferana, Abies balsamea, defoliation, fenitrothion; Quebec, sprayed by helicopter. 8, 300–305 (Smirnoff and Hardy)

barrier zone, traumatic tissue, wounds, *Liquidambar styraciflua*, decay, discolouration, compartmentalization; histochemistry, North Carolina U.S.A. 8, 389–397 (Moore)

Betula, electrical resistance, cankers, Endothia parasitica, Nectria galligena, Shigometer, Castanea; Massachusetts. 8, 162–167 (Sylvia and Tattar)

, fenitrothion, germination, Pinus banksiana, Picea glauca; toxicity. 8, 243-246 (Weinberger et al.)

Betula papyrifera, forest types, Picea mariana, Abies balsamea; Newfoundland, ribbed moraines. 8, 116–120 (Delaney and Cahill)

biomass, crown, Picea mariana; Alaska, foliage, cones, dead and live branches, bark and stem wood, prediction equations. 8, 36–41 (Barney et al.)

———, *Picea mariana*, forest type; subarctic Quebec, lichen woodlands, trees, shrubs, ground layer. **8**, 168–176 (Renez and Auclair)

———, whole-tree harvesting, nutrient depletion, *Pinus resinosa*, *Pinus banksiana*, *Pinus glauca*, *Populus tremuloides*, *Populus grandidentaia*; Minnesota, U.S.A., soils. **8**, 290–299 (Alban et al.)

browse, Alces alces, Abies balsamea, moose, regeneration; Quebec, Acer spicatum, Corylus cornuta. 8, 407–415 (Bédard et al.)

bud, apical dominance, morphogenesis, Picea sitchensis, plastochrone, primordia; Great Britain, cell dimensions. 8, 121–128 (Baxter and Cannell)

———, Larix decidua, leader, morphogenesis, Picea abies, Picea omorika, Picea sitchensis, tree form, phyllotaxy; Great Britain, lateral buds, position mechanism. 8, 129–137 (Cannell and Bowler)

budworm, fertilization, Abies balsamea, Choristoneura fumiferana; New Brunswick, budworm nutrition. 8, 364–374 (Shaw et al.)

cacodylic acid, Pinus nigra, interfascicular buds; proliferating buds. 8, 111-115 (Lheureux and Andre)

cankers, electrical resistance, Endothia parasitica, Nectria galligena, Shigometer, Castanea, Betula; Massachusetts. 8, 162—167 (Sylvia and Tattar)

Castanea, electrical resistance, cankers, Endothia parasitica, Nectria galligena, Shigometer, Betula; Massachusetts. 8, 162–167 (Sylvia and Tattar)

cation concentration, electrical resistance, Acer saccharum, Verticillium dahliae, decay; Massachusetts, U.S.A. 8, 322–327 (Malia and Tattar)

cation distribution, wounds, Acer rubrum; Maine, U.S.A., soil-applied potassium. 8, 228-231 (Blanchard et al.)

Choristoneura fumiferana, fertilization, budworm, Abies balsamea; New Brunswick, budworm nutrition. 8, 364–374 (Shaw et al.)

spruce budworm, *Abies balsamea*, defoliation, *Bacillus thuringiensis*, fenitrothion; Quebec, sprayed by helicopter, **8**, 300–305 (Smirnoff and Hardy)

clinal variation, *Picea mariana*, provenance, seed sources; tested in Ontario, initiation and cessation of growth, photoperiod and temperature. **8**, 463–473 (Morgenstern)

clonal variation, Populus, rabbits; Michigan, U.S.A., feeding selectivity. 8, 351-354 (Dickmann)

compartmentalization, traumatic tissue, barrier zone, wounds, Liquidambar styraciflua, decay, discolouration; histochemistry, North Carolina U.S.A. 8, 389–397 (Moore)

computer graphics, nursery stock; spatial mapping. 8, 261-263 (Lester and Martin)

Coniophora puteana, Armillaria mellea, Scytinostroma galactinum, Pinus banksiana, Pinus resinosa, Abies balsamea, root rotting; conifer saplings, northern Ontario. 8, 17–22 (Whitney and Myren)

container stock, *Pinus contorta*, wind throw, root morphology; mechanical instability, planting methods. **8**, 483–486 (Burdett)

containerized seedlings, irradiation, Picea mariana; greenhouse, growth chamber. 8, 316-321 (D'Aoust)

Pinus ponderosa, root form; field performance, 1.9-2 containers, 8, 232–236 (Owston and Seidel)

Cronartium comptoniae, Pinus banksiana, disease resistance, natural variation, provenance; tested in Minnesota, U.S.A. 8, 416–423 (Tauer)

——, sweetfern rust, *Pinus banksiana*; growth loss, cull, mortality, northern Ontario. **8**, 47–53 (Gross et al.) **crown**, biomass, *Picea mariana*; Alaska, foliage, cones, dead and live branches, bark and stem wood, prediction equations. **8**, 36–41 (Barney et al.)

cuttings, Salix; Alaska, taiga, vegetative propagation, five species. 8, 477-479 (Densmore and Zasada)

, vegetative propagation, *Pinus elliottii*, girdle; Mississippi, root-promoting chemicals, seasonal variation. **8**, 14–16 (Hare)

day length, *Pinus contorta*, genecology, temperature; wind-pollinated families, seedlings, greenhouse. **8**, 81–89 (Perry and Lotan)

decay, Acer saccharum, maple syrup, wound reaction; northeastern United States, succession of microorganisms, paraformaldehyde. 8, 54–60 (Walters and Shigo)

, electrical resistance, Acer saccharum, Verticillium dahliae, cation concentration; Massachusetts, U.S.A. 8, 322–327 (Malia and Tattar)

-, fungal mass, nitrogen-fixing bacteria, forest residues; Washington, U.S.A., glucosamine from amino acids. 8, 486-490 (Gurusiddaiah et al.)

-, traumatic tissue, barrier zone, wounds, Liquidambar styraciflua, discolouration, compartmentalization; histochemistry, North Carolina U.S.A. 8, 389-397 (Moore)

decay fungi, nitrogen fixation; Montana. 8, 341 - 345 (Larsen et al.)

decayed wood, Kraft pulping, Populus tremuloides; 8, 181-187 (Hunt et al.)

decomposition, Picea glauca, organic matter, humus; Alaska, cellulose bags, thinning, 8, 42-46 (Piene and Van Cleve)

defoliation, Quercus rubra, root regeneration; seedlings, spring shoot growth, dark periods, carbohydrates. 8, 67-72 (Larson)

spruce budworm, Choristoneura fumiferana, Abies balsamea, Bacillus thuringiensis, fenitrothion; Quebec, sprayed by helicopter. 8, 300-305 (Smirnoff and Hardy)

density, fertilization, Pinus resinosa; seedbeds, quality of planting stock. 8, 30-35 (Mullin and Bowdery)

discolouration, traumatic tissue, barrier zone, wounds, Liquidambar styraciflua, decay, compartmentalization; histochemistry, North Carolina U.S.A. 8, 389-397 (Moore)

disease resistance, Cronartium comptoniae, Pinus banksiana, natural variation, provenance; tested in Minnesota, U.S.A. 8, 416-423 (Tauer)

dormancy, dry matter accumulation, Picea sitchensis; Great Britain, winter activity, seedlings. 8, 207-213 (Bradbury and Malcolm)

drought, flower induction, lammas shoots, Pseudotsuga menziesii; 6- and 9-year grafts, France, seed cones. 8, 247-252 (Bonnet-Masimbert and Lanares)

dry matter accumulation, dormancy, Picea sitchensis; Great Britain, winter activity, seedlings. 8, 207-213 (Bradbury and Malcolm)

electrical resistance, Acer saccharum, Verticillium dahliae, cation concentration, decay; Massachusetts, U.S.A. 8, 322-327 (Malia and Tattar)

cankers, Endothia parasitica, Nectria galligena, Shigometer, Castanea, Betula; Massachusetts. 8, 162-167 (Sylvia and Tattar)

, Persea americana, Picea glauca, water potential; laboratory, New Brunswick, pressure bomb, psychrometer. 8, 73-80 (Dixon et al.)

-, phloem, Shigometer, Acer rubrum; field, Maine. 8, 90-93 (Carter and Blanchard)

emergence, temperature, flushing, heatsum, Pseudotsuga menziesii; modeling, nursery, growth chambers. 8, 23-29 (Bloomberg)

Endothia parasitica, electrical resistance, cankers, Nectria galligena, Shigometer, Castanea, Betula; Massachusetts. 8, 162-167 (Sylvia and Tattar)

energy balance, evapotranspiration; Quebec, stand of Abies balsamea. 8, 456-462 (McCaughey) evapotranspiration, energy balance; Quebec, stand of Abies balsamea. 8, 456-462 (McCaughey)

fenitrothion, germination, Pinus banksiana, Picea glauca, Betula; toxicity. 8, 243-246 (Weinberger et al.)

, seeds, Pinus strobus, germination; laboratory study, histochemistry, toxicology. 8, 155-161 (Weinberger et al.)

spruce budworm, Choristoneura fumiferana, Abies balsamea, defoliation, Bacillus thuringiensis; Quebec, sprayed by helicopter. 8, 300-305 (Smirnoff and Hardy)

fertilization, budworm, Abies balsamea, Choristoneura fumiferana; New Brunswick, budworm nutrition. 8, 364-374 (Shaw et al.)

-, density, Pinus resinosa; seedbeds, quality of planting stock. 8, 30-35 (Mullin and Bowdery)

organic layers, Picea mariana; urea, superphosphate, five seasons, New Brunswick. 8, 237-242, (Mahendrappa) fiber angle, spiral grain, heritability, radioisotope, Pinus nigra, Pinus pinaster; genetic prediction, juvenile-mature

correlation, France. 8, 280-289 (Arbez et al.)

fibre length, Fraxinus excelsior, internodal analysis; growth layer sequences. 8, 253-260 (Denne and Whitbread)

fire; Alberta, modeling, location allocation, airtankers. 8, 145-154 (Hodgson and Newstead)

, age-class; fire cycle, fire history. 8, 220-227 (Van Wagner)

fire ecology, fire intensity; discussion paper. 8, 491-492 (Van Wagner and Methven)

fire intensity, fire ecology; discussion paper. 8, 491-492 (Van Wagner and Methven)

flower induction, lammas shoots, drought, Pseudotsuga menziesii; 6- and 9-year grafts, France, seed cones. 8, 247-252 (Bonnet-Masimbert and Lanares)

flushing, emergence, temperature, heatsum, Pseudotsuga menziesii; modeling, nursery, growth chambers. 8. 23-29 (Bloomberg)

Fomes annosus, Armillaria mellea, infection centers, Pseudotsuga menziesii, Tsuga heterophylla; stumps, competition. 8, 177-180 (Morrison and Johnson)

forest openings, snow accumulation, meteorology; Alberta, ablation. 8, 380-388 (Golding and Swanson)

forest policies, sustained yield; British Columbia, forest industry communities, income, employment. 8, 61–66 (Byron)

forest residues, fungal mass, decay, nitrogen-fixing bacteria; Washington, U.S.A., glucosamine from amino acids. 8, 486–490 (Gurusiddaiah et al.)

forest type, biomass, *Picea mariana*; subarctic Quebec, lichen woodlands, trees, shrubs, ground layer. **8**, 168–176 (Rencz and Auclair)

forest types, Picea mariana, Betula papyrifera, Abies balsamea; Newfoundland, ribbed moraines. 8, 116–120 (Delaney and Cahill)

Fraxinus excelsior, fibre length, internodal analysis; growth layer sequences. 8, 253-260 (Denne and Whitbread)

frost damage, Juglans nigra, irrigation; Indiana, U.S.A. 8, 346-347 (Beineke)

fungal mass, decay, nitrogen-fixing bacteria, forest residues; Washington, U.S.A., glucosamine from amino acids. 8, 486–490 (Gurusiddaiah et al.)

genecology, Pinus contorta, day length, temperature; wind-pollinated families, seedlings, greenhouse. 8, 81–89 (Perry and Lotan)

germination, fenitrothion, Pinus banksiana, Picea glauca, Betula; toxicity. 8, 243-246 (Weinberger et al.)

et al.) seeds, fenitrothion, *Pinus strobus*; laboratory study, histochemistry, toxicology. **8**, 155–161 (Weinberger et al.)

girdle, vegetative propagation, Pinus elliottii, cuttings; Mississippi, root-promoting chemicals, seasonal variation. 8, 14–16 (Hare)

growth curves, Weibull function; height-age, volume-age. 8, 424-431 (Yang et al.)

heatsum, emergence, temperature, flushing, *Pseudotsuga menziesii*; modeling, nursery, growth chambers. **8**, 23–29 (Bloomberg)

heritability, spiral grain, fiber angle, radioisotope, Pinus nigra, Pinus pinaster; genetic prediction, juvenile-mature correlation, France. 8, 280-289 (Arbez et al.)

industry, sampling, pollution; British Columbia, grab and composite sampling, effluent. 8, 328-340 (Drechsler and Nemetz)

infection centers, Armillaria mellea, Fomes annosus, Pseudotsuga menziesii, Tsuga heterophylla; stumps, competition. 8, 177–180 (Morrison and Johnson)

infrared photography, Poria weirii, remote sensing; habitat types, colour photography, Idaho, U.S.A., spatial area. 8, 214—219 (Williams and Leaphart)

interfascicular buds, Pinus nigra, cacodylic acid; proliferating buds. 8, 111-115 (Lheureux and Andre)

internodal analysis, fibre length, Fraxinus excelsior; growth layer sequences. 8, 253–260 (Denne and Whitbread)

inventory, multiple use; decisions, sampling error, Virginia, modeling. 8, 100-110 (Burkhart et al.)

irradiance, Populus euramerica; rooted cuttings, infrared gas analyzer, three clones, growth, photosynthesis. 8, 94–99 (Fasehun)

irradiation, Picea mariana, containerized seedlings; greenhouse, growth chamber. 8, 316-321 (D'Aoust)

irrigation, Juglans nigra, frost damage; Indiana, U.S.A. 8, 346-347 (Beineke)

isolation, Armillaria mellea; selective media. 8, 348-350 (Whitney et al.)

Juglans nigra, frost damage, irrigation; Indiana, U.S.A. 8, 346-347 (Beineke)

Kraft pulping, Populus tremuloides, decayed wood; 8, 181-187 (Hunt et al.)

lammas shoots, flower induction, drought, *Pseudotsuga menziesii*; 6- and 9-year grafts, France, seed cones. 8, 247–252 (Bonnet-Masimbert and Lanares)

Larix, phyllotaxy, Picea; computer simulation, primordium, lateral buds, Fibonacci number. 8, 138-141 (Cannell and Bowler)

Larix decidua, bud, leader, morphogenesis, Picea abies, Picea omorika, Picea sitchensis, tree form, phyllotaxy; Great Britain, lateral buds, position mechanism. 8, 129–137 (Cannell and Bowler)

leader, bud, Larix decidua, morphogenesis, Picea abies, Picea omorika, Picea sitchensis, tree form, phyllotaxy; Great Britain, lateral buds, position mechanism. 8, 129–137 (Cannell and Bowler)

Liquidambar styraciflua, traumatic tissue, barrier zone, wounds, decay, discolouration, compartmentalization; histochemistry, North Carolina U.S.A. 8, 389–397 (Moore)

maple syrup, Acer saccharum, wound reaction, decay; northeastern United States, succession of microorganisms, paraformaldehyde. 8, 54–60 (Walters and Shigo)

meteorology, snow accumulation, forest openings; Alberta, ablation. 8, 380-388 (Golding and Swanson)

mineralization, spacing, Abies bulsamea, organic layers, temperature; Nova Scotia. 8, 398-406 (Piene)

moisture stress, *Pinus contorta*, osmotic potentials, water potential; Idaho, U.S.A., needles and stem tissues. **8**, 439–441 (Spomer and Stoszek)

moose, Alces alces, Abies balsamea, browse, regeneration; Quebec, Acer spicatum, Corylus cornuta. 8, 407-415 (Bédard et al.)

morphogenesis, apical dominance, bud, *Picea sitchensis*, plastochrone, primordia; Great Britain, cell dimensions. **8**, 121–128 (Baxter and Cannell)

———, bud, Larix decidua, leader, Picea abies, Picea omorika, Picea sitchensis, tree form, phyllotaxy; Great Britain, lateral buds, position mechanism. 8, 129–137 (Cannell and Bowler)

multiple use, inventory; decisions, sampling error, Virginia, modeling. 8, 100-110 (Burkhart et al.)

mycorrhizae, Acer saccharum; Massachusetts, maple decline. 8, 375-379 (Spitko et al.)

natural variation, Cronartium comptoniae, Pinus banksiana, disease resistance, provenance; tested in Minnesota, U.S.A. 8, 416-423 (Tauer)

Nectria galligena, electrical resistance, cankers, Endothia parasitica, Shigometer, Castanea, Betula; Massachusetts. 8, 162—167 (Sylvia and Tattar)

needles, temperature, *Pinus silvestris*; Sweden, ultraminiature thermocouple. **8**, 480–482 (Christersson and Sandstedt) **nitrogen availability**, whole-tree harvesting, organic layers; modeling, forest floor biomass. **8**, 306–315 (Aber et al.)

nitrogen fixation, decay fungi; Montana. 8, 341-345 (Larsen et al.)

nitrogen-fixing bacteria, fungal mass, decay, forest residues; Washington, U.S.A., glucosamine from amino acids. 8, 486–490 (Gurusiddaiah et al.)

nursery stock, computer graphics; spatial mapping. 8, 261-263 (Lester and Martin)

nutrient depletion, whole-tree harvesting, Pinus resinosa, Pinus banksiana, Pinus glauca, Populus tremuloides, Populus grandidentata, biomass; Minnesota, U.S.A., soils. 8, 290–299 (Alban et al.)

nutrient losses, *Picea glauca*, *Abies lasiocarpa*, roots; British Columbia, management, logging, fine-root biomass, mineral elements. **8**, 265–279 (Kimmins and Hawkes)

Persea americana, Picea glauca, electrical resistance, water potential; laboratory, New Brunswick, pressure bomb, psychrometer. 8, 73–80 (Dixon et al.)

phloem, electrical resistance, Shigometer, Acer rubrum; field, Maine. 8, 90-93 (Carter and Blanchard)

and Bowler)

Picea, phyllotaxy, Larix; computer simulation, primordium, lateral buds, Fibonacci number. 8, 138-141 (Cannell and Bowler)

Picea abies, bud, Larix decidua, leader, morphogenesis, Picea omorika, Picea sitchensis, tree form, phyllotaxy; Great Britain, lateral buds, position mechanism. 8, 129–137 (Cannell and Bowler)

Picea glauca, Abies lasiocarpa, roots, nutrient losses; British Columbia, management, logging, fine-root biomass, mineral

elements. 8, 265–279 (Kimmins and Hawkes)

, fenitrothion, germination, Pinus banksiana, Betula; toxicity. 8, 243-246 (Weinberger et al.)

organic matter, humus, decomposition; Alaska, cellulose bags, thinning. 8, 42-46 (Piene and Van Cleve)

——, Persea americana, electrical resistance, water potential; laboratory, New Brunswick, pressure bomb, psychrometer, 8, 73—80 (Dixon et al.)

Picea mariana, biomass, crown; Alaska, foliage, cones, dead and live branches, bark and stem wood, prediction equations. 8, 36–41 (Barney et al.)

—, biomass, forest type; subarctic Quebec, lichen woodlands, trees, shrubs, ground layer. **8**, 168–176 (Rencz and Auclair)

, forest types, Betula papyrifera, Abies balsamea; Newfoundland, ribbed moraines. 8, 116-120 (Delaney and Cahill)

, irradiation, containerized seedlings; greenhouse, growth chamber. 8, 316-321 (D'Aoust)

, provenance, seed sources, clinal variation; tested in Ontario, initiation and cessation of growth, photoperiod and temperature. 8, 463–473 (Morgenstern)

Picea omorika, bud, *Larix decidua*, leader, morphogenesis, *Picea abies, Picea sitchensis*, tree form, phyllotaxy; Great Britain, lateral buds, position mechanism. 8, 129–137 (Cannell and Bowler)

Picea sitchensis, apical dominance, bud, morphogenesis, plastochrone, primordia; Great Britain, cell dimensions. 8, 121–128 (Baxter and Cannell)

—, bud, Larix decidua, leader, morphogenesis, Picea abies, Picea omorika, tree form, phyllotaxy; Great Britain, lateral buds, position mechanism. 8, 129–137 (Cannell and Bowler)

———, dormancy, dry matter accumulation; Great Britain, winter activity, seedlings. 8, 207–213 (Bradbury and Malcolm)

Pinus, principal component, stem form, taper; southeastern United States. 8, 188–197 (Liu and Keister)

Pinus banksiana, Armillaria mellea, Scytinostroma galactinum, Coniophora puteana, Pinus resinosa, Abies balsamea, root rotting; conifer saplings, northern Ontario. 8, 17–22 (Whitney and Myren)

——, Cronartium comptoniae, disease resistance, natural variation, provenance; tested in Minnesota, U.S.A. 8, 416–423 (Tauer)

, fenitrothion, germination, Picea glauca, Betula; toxicity. 8, 243-246 (Weinberger et al.)

———, whole-tree harvesting, nutrient depletion, *Pinus resinosa*, *Pinus glauca*, *Populus tremuloides*, *Populus grandidentata*, biomass; Minnesota, U.S.A., soils. **8**, 290–299 (Alban et al.)

Pinus contortu, container stock, wind throw, root morphology; mechanical instability, planting methods. 8, 483-486 (Burdett)

———, genecology, day length, temperature; wind-pollinated families, seedlings, greenhouse. **8**, 81–89 (Perry and Lotan)

moisture stress, osmotic potentials, water potential; Idaho, U.S.A., needles and stem tissues. 8, 439–441 (Spomer and Stoszek)

Pinus elliottii, vegetative propagation, girdle, cuttings; Mississippi, root-promoting chemicals, seasonal variation. 8, 14–16 (Hare)

Pinus glauca, whole-tree harvesting, nutrient depletion, Pinus resinosa, Pinus banksiana, Populus tremuloides, Populus grandidentata, biomass; Minnesota, U.S.A., soils. 8, 290–299 (Alban et al.)

Pinus nigra, cacodylic acid, interfascicular buds; proliferating buds. 8, 111-115 (Lheureux and Andre)

- spiral grain, fiber angle, heritability, radioisotope, *Pinus pinaster*; genetic prediction, juvenile-mature correlation, France. **8**, 280–289 (Arbez et al.)

Pinus pinaster, spiral grain, fiber angle, heritability, radioisotope, Pinus nigra; genetic prediction, juvenile-mature correlation, France. 8, 280-289 (Arbez et al.)
Pinus ponderosa, containerized seedlings, root form; field performance, 1.9-8 containers. 8, 232-236 (Owston and

Pinus ponderosa, containerized seedlings, root form; field performance, 1.9-8 containers, 8, 232–236 (Owston and Seidel)

Pinus resinosa, Armillaria mellea, Scytinostroma galactinum, Coniophora puteana, Pinus banksiana, Abies balsamea, root rotting; conifer saplings, northern Ontario. 8, 17–22 (Whitney and Myren)

, fertilization, density; seedbeds, quality of planting stock. 8, 30-35 (Mullin and Bowdery)

———, whole-tree harvesting, nutrient depletion, *Pinus banksiana*, *Pinus glauca*, *Populus tremuloides*, *Populus grandidentata*, biomass; Minnesota, U.S.A., soils. **8**, 290–299 (Alban et al.) *Pinus silvestris*, temperature, needles; Sweden, ultraminiature thermocouple. **8**, 480–482 (Christersson and Sandstedt)

Pinus strobus, pollutants, wax; New Brunswick, scanning electron microscopy. **8**, 474–477 (Percy and Riding)

et al.) , seeds, fenitrothion, germination; laboratory study, histochemistry, toxicology. 8, 155-161 (Weinberger et al.)

Weibull function, seed dispersal; hardwood stands, modeling, seedling spatial patterns. 8, 10-13 (Rudis et al.)

plastochrone, apical dominance, bud, morphogenesis, Picea sitchensis, primordia; Great Britain, cell dimensions. 8, 121–128 (Baxter and Cannell)

pollutants, Pinus strobus, wax; New Brunswick, scanning electron microscopy. 8, 474-477 (Percy and Riding)

pollution, Acer saccharum; Ohio River Valley, U.S.A., airborne, seven stands, richness and deversity. 8, 432–438 (McClenahen)

and Nemetz) ——, sampling, industry; British Columbia, grab and composite sampling, effluent. **8**, 328–340 (Drechsler and Nemetz)

Populus, rabbits, clonal variation; Michigan, U.S.A., feeding selectivity. 8, 351-354 (Dickmann)

Populus euramerica, irradiance; rooted cuttings, infrared gas analyzer, three clones, growth, photosynthesis. 8, 94–99 (Fasehun)

Populus grandidentata, whole-tree harvesting, nutrient depletion, Pinus resinosa, Pinus banksiana, Pinus glauca, Populus tremuloides, biomass; Minnesota, U.S.A., soils. 8, 290-299 (Alban et al.)

Populus tremuloides, Kraft pulping, decayed wood; 8, 181-187 (Hunt et al.)

whole-tree harvesting, nutrient depletion, Pinus resinosa, Pinus banksiana, Pinus glauca, Populus grandidentata, biomass; Minnesota, U.S.A., soils. 8, 290-299 (Alban et al.) Poria weirii, infrared photography, remote sensing; habitat types, colour photography, Idaho, U.S.A., spatial area. 8, 214

219 (Williams and Leaphart)

primordia, apical dominance, bud, morphogenesis, Picea sitchensis, plastochrone; Great Britain, cell dimensions. 8, 121-128 (Baxter and Cannell)

principal component, stem form, taper, Pinus; southeastern United States, 8, 188-197 (Liu and Keister)

provenance, Cronartium comptoniae, Pinus banksiana, disease resistance, natural variation; tested in Minnesota, U.S.A. 8, 416-423 (Tauer)

Picea mariana, seed sources, clinal variation; tested in Ontario, initiation and cessation of growth, photoperiod and temperature. 8, 463-473 (Morgenstern) Pseudotsuga menziesii, Armillaria mellea, infection centers, Fomes annosus, Tsuga heterophylla; stumps, competition, 8,

177-180 (Morrison and Johnson)

-, emergence, temperature, flushing, heatsum; modeling, nursery, growth chambers. 8, 23-29 (Bloomberg) flower induction, lammas shoots, drought; 6- and 9-year grafts, France, seed cones. 8, 247-252 (Bonnet-Masimbert and Lanares)

Quercus rubra, defoliation, root regeneration; seedlings, spring shoot growth, dark periods, carbohydrates. 8, 67-72

rabbits, Populus, clonal variation; Michigan, U.S.A., feeding selectivity. 8, 351-354 (Dickmann)

radiation balance, reflection coefficient; Quebec, stand of Abies balsamea, logged. 8, 450-455 (McCaughey)

radioisotope, spiral grain, fiber angle, heritability, Pinus nigra, Pinus pinaster, genetic prediction, juvenile-mature correlation, France. 8, 280-289 (Arbez et al.)

reflection coefficient, radiation balance; Quebec, stand of Abies balsamea, logged, 8, 450-455 (McCaughey)

regeneration, Alces alces, Abies balsamea, moose, browse; Quebec, Acer spicatum, Corylus cornuta. 8, 407-415 (Bédard et al.)

remote sensing, Poria weirii, infrared photography; habitat types, colour photography, Idaho, U.S.A., spatial area. 8, 214-219 (Williams and Leaphart)

root; landslides, shear and tensile strength, deterioration, California. 8, 142–144 (Ziemer)

root form, containerized seedlings, Pinus ponderosa; field performance, 1.9-8 containers. 8, 232-236 (Owston and

root morphology, container stock, Pinus contorta, wind throw; mechanical instability, planting methods. 8, 483-486 (Burdett)

root regeneration, Quercus rubra, defoliation; seedlings, spring shoot growth, dark periods, carbohydrates. 8, 67-72 (Larson)

root rotting, Armillaria mellea, Scytinostroma galactinum, Coniophora puteana, Pinus banksiana, Pinus resinosa, Abies balsamea; conifer saplings, northern Ontario. 8, 17-22 (Whitney and Myren)

roots, Picea glauca, Abies lusiocarpa, nutrient losses; British Columbia, management, logging, fine-root biomass, mineral elements. 8, 265-279 (Kimmins and Hawkes)

Salix, cuttings; Alaska, taiga, vegetative propagation, five species. 8, 477-479 (Densmore and Zasada)

sampling, industry, pollution; British Columbia, grab and composite sampling, effluent. 8, 328-340 (Drechsler and Nemetz)

Scytinostroma galactinum, Armillaria mellea, Coniophora puteana, Pinus banksiana, Pinus resinosa, Abies balsamea, root rotting; conifer saplings, northern Ontario. 8, 17-22 (Whitney and Myren) seed dispersal, Weibull function, Pinus strobus; hardwood stands, modeling, seedling spatial patterns. 8, 10-13 (Rudis et

al.) seed sources, Picea mariana, provenance, clinal variation; tested in Ontario, initiation and cessation of growth,

photoperiod and temperature. 8, 463-473 (Morgenstern) seeds, fenitrothion, Pinus strobus, germination; laboratory study, histochemistry, toxicology. 8, 155-161 (Weinberger et

Shigometer, electrical resistance, cankers, Endothia parasitica, Nectria galligena, Castanea, Betula; Massachusetts. 8, 162-167 (Sylvia and Tattar)

electrical resistance, phloem, Acer rubrum; field, Maine. 8, 90-93 (Carter and Blanchard) snow accumulation, forest openings, meteorology; Alberta, ablation. 8, 380-388 (Golding and Swanson)

spacing, mineralization, Abies balsamea, organic layers, temperature; Nova Scotia. 8, 398-406 (Piene)

- spiral grain, fiber angle, heritability, radioisotope, Pinus nigra, Pinus pinaster; genetic prediction, juvenile-mature correlation, France. 8, 280-289 (Arbez et al.)
- spruce budworm, Choristoneura fumiferana, Abies balsamea, defoliation, Bacillus thuringiensis, fenitrothion; Quebec, sprayed by helicopter. 8, 300–305 (Smirnoff and Hardy)
- stem form, principal component, taper, Pinus; southeastern United States. 8, 188-197 (Liu and Keister)
- sustained yield, forest policies; British Columbia, forest industry communities, income, employment. 8, 61-66 (Byron)
- sweetfern rust, Cronartium comptoniae, Pinus banksiana; growth loss, cull, mortality, northern Ontario. 8, 47-53 (Gross et al.)
- taper, principal component, stem form, Pinus; southeastern United States. 8, 188-197 (Liu and Keister)
- temperature, emergence, flushing, heatsum, *Pseudotsuga menziesii*; modeling, nursery, growth chambers. **8**, 23–29 (Bloomberg)
- needles, *Pinus silvestris*; Sweden, ultraminiature thermocouple. **8**, 480–482 (Christersson and Sandstedt)

 , *Pinus contorta*, genecology, day length; wind-pollinated families, seedlings, greenhouse. **8**, 81–89 (Perry and Lotan)
- spacing, mineralization, Abies balsamea, organic layers; Nova Scotia. 8, 398-406 (Piene)
- traumatic tissue, barrier zone, wounds, Liquidambar styraciflua, decay, discolouration, compartmentalization; histochemistry, North Carolina U.S.A. 8, 389–397 (Moore)
- tree form, bud, Larix decidua, leader, morphogenesis, Picea abies, Picea omorika, Picea sitchensis, phyllotaxy; Great Britain, lateral buds, position mechanism. 8, 129–137 (Cannell and Bowler)
- Tsuga heterophylla, Armillaria mellea, infection centers, Fomes annosus, Pseudotsuga menziesii; stumps, competition. 8, 177–180 (Morrison and Johnson)
- vegetative propagation, *Pinus elliottii*, girdle, cuttings; Mississippi, root-promoting chemicals, seasonal variation. **8**, 14–16 (Hare)
- Verticillium dahliae, electrical resistance, Acer saccharum, cation concentration, decay; Massachusetts, U.S.A. 8, 322–327 (Malia and Tattar)
- water potential. Persea americana, Picea glauca, electrical resistance; laboratory, New Brunswick, pressure bomb, psychrometer. 8, 73–80 (Dixon et al.)
- wax, *Pinus strobus*, pollutants; New Brunswick, scanning electron microscopy. **8**, 474–477 (Percy and Riding)
- Weibull function. growth curves; height-age, volume-age. 8, 424-431 (Yang et al.)
- Pinus strobus, seed dispersal; hardwood stands, modeling, seedling spatial patterns. 8, 10-13 (Rudis et
- whole-tree harvesting, nitrogen availability, organic layers; modeling, forest floor biomass. 8, 306–315 (Aber et al.)
- wind throw, container stock, *Pinus contorta*, root morphology; mechanical instability, planting methods. **8**, 483–486 (Burdett)
- wound reaction, Acer saccharum, maple syrup, decay; northeastern United States, succession of microorganisms, paraformaldehyde. 8, 54–60 (Walters and Shigo)
- wounds, cation distribution, Acer rubrum; Maine, U.S.A., soil-applied potassium. 8, 228-231 (Blanchard et al.)
- , traumatic tissue, barrier zone, *Liquidambar styraciftua*, decay, discolouration, compartmentalization; histochemistry, North Carolina U.S.A. 8, 389–397 (Moore)

X-ray densitometry, annual ring; radial and transverse sections. 8, 442-449 (Cown and Parker)

